

COMT Val¹⁵⁸Met genotype is associated with reward learning: a replication study and meta-analysis

N. S. Corral-Frías^{†,‡,*}, D. A. Pizzagalli[§],
J. M. Carré[¶], L. J. Michalski[‡], Y. S. Nikolova^{**},
R. H. Perlis^{††,‡‡}, J. Fagerness^{‡‡}, M. R. Lee^{§§},
E. Drabant Conley^{¶¶}, T. M. Lancaster^{***},
S. Haddad^{‡‡}, A. Wolf^{†††}, J. W. Smoller^{††,‡‡},
A. R. Hariri^{‡‡‡} and R. Bogdan^{‡,§§§}

[†]Psychiatry Department, [‡]BRAIN Laboratory, Department of Psychology, Washington University in St. Louis, St. Louis, MO, USA, [§]Center For Depression, Anxiety and Stress Research and Neuroimaging Center, McLean Hospital and Harvard Medical School, Belmont, MA, USA, [¶]Nipissing University, North Bay, Ontario, Canada, ^{**}Centre for Addiction and Mental Health Toronto, Ontario Canada, ^{††}Massachusetts General Hospital and Harvard Medical School, Cambridge, MA, USA, ^{‡‡}Psychiatric and Neurodevelopmental Genetics Unit, Center for Human Genetic Research, Massachusetts General Hospital, Boston, MA, USA, ^{§§}National Institute on Drug Abuse, Intramural Research Program, Baltimore, MD, USA, ^{¶¶}23 and Me, ^{***}Neuroscience and Mental Health Research Institute, Cardiff University, Cardiff, UK, ^{†††}Department of Psychiatry Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA, ^{‡‡‡}Laboratory of NeuroGenetics, Department of Psychology and Neuroscience, Duke University, Durham, NC, USA, and ^{§§§}Division of Biology and Biomedical Sciences, Washington University in St. Louis, St. Louis, MO, USA

*Corresponding authors: N. S. Corral-Frías and R. Bogdan, Campus Box 1125, Psychology Bldg., Washington University in St. Louis, One Brookings Dr., St. Louis, MO 63130, USA. E-mail: corraln@psychiatry.wustl.edu and rbogdan@wustl.edu

Identifying mechanisms through which individual differences in reward learning emerge offers an opportunity to understand both a fundamental form of adaptive responding as well as etiological pathways through which aberrant reward learning may contribute to maladaptive behaviors and psychopathology. One candidate mechanism through which individual differences in reward learning may emerge is variability in dopaminergic reinforcement signaling. A common functional polymorphism within the catechol-O-methyl transferase gene (COMT; rs4680, Val¹⁵⁸Met) has been linked to reward learning, where homozygosity for the Met allele (linked to heightened prefrontal dopamine function and decreased dopamine synthesis in the midbrain) has been associated with relatively increased reward learning. Here, we used a probabilistic reward learning task to assess response bias, a behavioral form of reward learning, across three separate samples that were combined for analyses (age: 21.80 ± 3.95; n = 392;

268 female; European-American: n = 208). We replicate prior reports that COMT rs4680 Met allele homozygosity is associated with increased reward learning in European-American participants ($\beta = 0.20$, $t = 2.75$, $P < 0.01$; $\Delta R^2 = 0.04$). Moreover, a meta-analysis of 4 studies, including the current one, confirmed the association between COMT rs4680 genotype and reward learning (95% CI -0.11 to -0.03 ; $z = 3.2$; $P < 0.01$). These results suggest that variability in dopamine signaling associated with COMT rs4680 influences individual differences in reward which may potentially contribute to psychopathology characterized by reward dysfunction.

Keywords: Anhedonia, COMT, dopamine, meta-analysis, response bias, reward

Received 13 January 2016, revised 25 March 2016, accepted for publication 14 April 2016

Blunted reward processing is a cardinal feature of depression that is frequently observed across other forms of psychopathology, including posttraumatic stress disorder (PTSD), schizophrenia and substance use disorders (Garfield *et al.* 2014; Gorwood 2008; Hatzigiakoumis *et al.* 2011; Pizzagalli 2014). Consistent with theoretical speculation (Loas 1996; Meehl 1975), emerging evidence suggests that diminished hedonic capacity may provide trait-like vulnerability to psychopathology (Corral-Frías *et al.* 2015; Nikolova *et al.* 2012), making it important to identify the origin and mechanisms underlying individual differences in reward processing. Guided by evidence that variability in reward processing is heritable (Bogdan & Pizzagalli 2009; Wichers *et al.* 2007) and linked to dopaminergic (DA) system function (Schultz 2015), genetic association studies of reward processing have focused primarily on functional polymorphisms within DA-related proteins (Bogdan *et al.* 2013; Forbes *et al.* 2009; Nikolova *et al.* 2011).

The most studied DA-related polymorphism in psychiatric and behavioral genetics to date is rs4680 (Val¹⁵⁸Met) within the catechol-O-methyl transferase (COMT) gene (COMT) (Buckholtz & Meyer-Lindenberg 2012; Gatt *et al.* 2015), which codes for a catabolic catecholamine enzyme (Mannisto & Kaakkola 1999). Along with the DA transporter (DAT), the COMT enzyme is one of the primary synaptic regulators of DA. Unlike DAT, which is primarily expressed in subcortical regions, COMT is widely expressed in the prefrontal cortex (PFC) and is the primary constraint of prefrontal synaptic DA transmission (Tunbridge *et al.* 2004). Met (A) allele homozygosity at rs4680 is associated with a 40% reduction in COMT activity relative to Val (G) allele homozygosity (Chen *et al.* 2004).

This genotype-dependent reduction in COMT activity results in relatively higher DA levels in the PFC (Meyer-Lindenberg *et al.* 2005; Slifstein *et al.* 2008). While the direct effect of *COMT* Val¹⁵⁸Met genotype is primarily on cortical DA, there is evidence that such cortical effects may indirectly modulate subcortical DA signaling (Akil *et al.* 2003; Meyer-Lindenberg *et al.* 2005; Scornaiencki *et al.* 2009; Seamans & Yang 2004); indeed, the Met allele has been associated with decreased midbrain DA synthesis (Akil *et al.* 2003; Meyer-Lindenberg *et al.* 2005), which may facilitate the detection of phasic DA shifts critical for reward prediction errors and reinforcement learning (Bilder *et al.* 2004; Bogdan *et al.* 2011; Santesso *et al.* 2008; Schultz 2002, 2007). Consistent with this notion, the Met allele has been linked to heightened behavioral reward learning (Lancaster *et al.* 2012, 2015), elevated positive affect in response to reward (Wichers *et al.* 2007), and reward-seeking behavior (Lancaster *et al.* 2012) as well as reduced anhedonic symptoms in relatives of schizophrenia patients (Docherty & Sponheim 2008). Such individual differences in reward-related behavior may underlie associations between *COMT* rs4680 genotype and psychopathology (Antypa *et al.* 2013; Bogdan *et al.* 2013).

Given emergent evidence linking *COMT* genotype to reward learning (Frank *et al.* 2007; Lancaster *et al.* 2012, 2015), and recent concerns of lack replication in behavioral genetics (Duncan & Keller 2011; Plomin *et al.* 2016) the present study examined whether *COMT* genotype (rs4680) is associated with reward learning using data from three samples. Based on prior research (Lancaster *et al.* 2012, 2015), we hypothesized that individuals homozygous for the low activity Met allele would have increased reward learning (i.e. greater response bias to more rewarded cues). Lastly, we conducted a meta-analysis of published studies examining associations between *COMT* rs4680 genotype and behavioral reward learning as measured by a probabilistic reward learning task.

Materials and methods

Participants

Participants ($n=392$) were recruited for three independent studies from the general and college community in the greater Boston, Massachusetts (Samples 1–2) and Durham, North Carolina (Sample 3) areas. Following quality control within each sample described below, the final total sample included 303 participants [age: 21.80 ± 3.95 ; 209 (69%) female; ethnicity: 208 (68.6%) European/European American, 33 (10.9%) African/African-American, 39 (12.9%) Asian/Asian-American, 10 (3.3%) Hispanic, 11 (3.6%) multiracial or other, 2 did not report (0.7%); Table S1, Supporting Information]. Because the relationship between this polymorphism and response bias has only been characterized in European and European-American samples (Goetz *et al.* 2013; Lancaster *et al.* 2012, 2015), and owing to evidence for differential associations across ancestral origins in other phenotypes (Lee & Prescott 2014), primary analyses were conducted on European-American participants (Table 1) with supplemental analyses conducted in the entire sample (Appendix S1). All subjects gave written informed consent and studies were approved by the Harvard University and Duke University Institutional Review Boards.

Sample 1

Healthy female participants ($n=84$) aged 18–25 were recruited from the greater Boston community. Exclusionary criteria included

Table 1: Sample demographics for composite European American sample

	Sample 1 ($n=56$)			Sample 2 ($n=119$)			Sample 3 ($n=33$)			All Samples ($n=208$)		
	Met/Met ($n=14$)	Val/Val, Val/Met ($n=42$)	Test statistic+	Met/Met ($n=30$)	Val/Val, Val/Met ($n=89$)	COMT relation	Met/Met ($n=11$)	Val/Val, Val/Met ($n=22$)	Test statistic+	Met/Met ($n=55$)	Val/Val, Val/Met ($n=153$)	Test statistic+
Age (SD)	21.8 ± 2.22	21.9 ± 1.63	$t=0.18$ $P=0.85$	21.5 ± 3.00	23.4 ± 4.73	$t=2.10$ $P=0.038^*$	19.0 ± 1.09	19.6 ± 1.21	$t=1.46$ $P=0.15$	21.0 ± 2.74	22.5 ± 4.08	$t=2.407$ $P=0.017^*$
Sex (% female)	14 (100%)	42 (100%)	n/a	13 (43.33%)	48 (53.93%)	$\chi^2=1.00$ $P=0.315$	5 (45.45%)	13 (59.09%)	$\chi^2=0.55$ $P=0.458$	32 (58.18%)	103 (67.32%)	$\chi^2=1.48$ $P=0.22$

* $P < 0.05$.

+ Comparison between Met homozygotes and Val carriers.

left-handedness, color blindness, past or present neurological, psychiatric, hormonal, or metabolic disturbances, and self-report ethnicity (i.e. only participants with two parents of European ancestry were included). Participants provided written informed consent to a protocol approved by the Committee on the Use of Human Subjects in Research at Harvard University and received either course credit or \$10/h as well as additional compensation earned (\$15) during the probabilistic reward learning task (described below). Data were excluded from analyses for the following reasons: genotyping was not conducted ($n=19$), task non-compliance (i.e. predominantly pressing only one button) or below chance accuracy ($n=6$), technical difficulties (i.e. equipment did not function properly; $n=3$) and failed genotyping ($n=1$), leaving a final sample of 56 for analyses. Three prior manuscripts have been published using these data evaluating reinforcement learning parameters (Huys *et al.* 2013) and associations between stress and genetic variation in the hypothalamic-pituitary-adrenal axis (Bogdan *et al.* 2010, 2011).

Sample 2

Participants ($n=214$; 123 female) aged 18–64 were recruited from Harvard University and the greater Boston community. Exclusionary criteria included current medical illness, attention-deficit hyperactivity disorder (ADHD), head injury, loss of consciousness, seizures, current alcohol/substance abuse or dependence, smoking, use of psychotropic medications during the last 2 weeks, pregnancy, or left handedness. Participants provided written informed consent to a protocol approved by the Committee on the Use of Human Subjects in Research at Harvard University and received course credit or \$5 for participation and won additional money (average \$6.00; between \$5.80 and \$6.20) while completing the reward task. Collected data were excluded ($n=41$) from analyses due to task noncompliance (i.e. predominantly pressing only one button, below chance accuracy, or an inadequate reward ratio exposure, $n=36$), as well as failed genotyping ($n=5$) leaving a final sample of 173 (105 female) participants (119 European American; 61 female) for the present analyses. Three prior manuscripts have been published using these data evaluating computation reinforcement learning parameters (Huys *et al.* 2013) associations between stress and genetic variation within the HPA axis (Bogdan *et al.* 2010) and neural substrates of reward learning (Santesso *et al.* 2008).

Sample 3

A subset of participants ($n=108$; 70 females) enrolled in the ongoing Duke Neurogenetics Study (DNS; Carey *et al.* 2015; Corral-Frias *et al.* 2015; Nikolova *et al.* 2014) completed the probabilistic reward learning task described below. Participants provided written informed consent to a protocol approved by Duke University and received \$5 for their time and won an additional \$5 while completing the task. Study exclusion criteria included: medical diagnoses of cancer, stroke, diabetes requiring insulin treatment, chronic kidney or liver disease, or lifetime history of psychotic symptoms; use of psychotropic, glucocorticoid, or hypolipidemic medication; and/or conditions affecting cerebral blood flow and metabolism (e.g. hypertension). As the DNS seeks to establish broad variability in multiple behavioral phenotypes related to psychopathology, diagnosis of current DSM-IV Axis I and select Axis II disorders (Antisocial Personality Disorder and Borderline Personality Disorder) were not exclusionary. Collected data were excluded from analyses due to task noncompliance (e.g. pressing one button exclusively, $n=6$) or because genotyping was not conducted ($n=26$). The final sample included a total of 74 (48 female) participants (33 European-American, 18 female).

Reward learning task and data processing

The computer task, which was adapted from prior studies (Pizzagalli *et al.* 2005; Tripp & Alsup 1999), was presented on a PC using E-prime software (Psychology Software Tools, Inc, Pittsburgh, PA, USA). Notably, reward learning as measured by this task is (1) heritable (Bogdan & Pizzagalli 2009), (2) associated with depression and anhedonia (Luking *et al.* 2015a,b; Pizzagalli *et al.* 2005), (3) linked to treatment outcomes (Vrieze *et al.* 2013) and smoking behaviors in depressed patients (Liverant *et al.* 2014), (4) associated




Figure 1: Schematic diagram of the reward learning task.

Participants are instructed to press a button on the keyboard to indicate whether a long or short mouth is presented (100 milliseconds) within a schematic face. Following *some*, but not all correct responses, participants received a monetary reward of 5 cents. One stimulus (rich) was rewarded three times more than the other (lean). In Sample 1, two different stimuli were used: mouth and nose. In this sample participants had to indicate whether a long (mouth, 11.00 mm; nose, 5.31 mm) or short (mouth, 10.00 mm; nose, 5.00 mm) stimulus was presented. In Sample 2, only mouth stimuli (long mouth: 13 mm, short mouth: 11.5 mm) were presented. In Sample 3, only mouth stimuli were presented, and these were the same length as in Sample 1. Figure adapted from Pizzagalli *et al.* (2005).

with depression resistance among anxious individuals (Morris & Rottenberg 2015), (5) linked to reward-related striatal function (Santesso *et al.* 2008) and DA release (Vrieze *et al.* 2013) and (6) blunted by stress (Bogdan & Pizzagalli 2006; Bogdan *et al.* 2011) and nicotine withdrawal (Pergadia *et al.* 2014). Briefly, participants are instructed to press a button on a button box or a keyboard to indicate whether a long or short mouth or nose is presented (100 milliseconds) within a schematic face (see Fig. 1). Importantly, the small size difference between stimuli and brief exposure time makes it difficult to discern which stimulus is presented. Participants are told that *some*, but not all correct responses, will result in correct feedback and a monetary reward. One of the stimuli (i.e. either long or short), the 'rich' stimulus, is rewarded three times more frequently than the other, 'lean' stimulus (stimulus types and buttons were counterbalanced across participants). Under these contingencies, humans and non-human animals develop a response bias for the more frequently rewarded, 'rich' stimulus (Der-Avakian *et al.* 2013; Herrnstein 1961; Lauwereyns *et al.* 2002; Pizzagalli *et al.* 2005; Tripp & Alsup 1999).

The task consists of three blocks with 40% of trials per block receiving a reward (task in Samples 1 and 3 consisted of 80 blocks whereas in Sample 2 it consisted of 100 blocks; accordingly, 32 of the trials in Samples 1 and 3 were rewarded and 40 trials in Sample 2 were rewarded). The 'rich' and 'lean' stimuli were presented with equal frequency, but, unknown to the participants, the reward feedback is asymmetrical in favor of the 'rich' stimulus (3 'rich' to 1 'lean' reward ratio; participants in Samples 1 and 3 received reward for 24 and 8 of the rich and lean stimulus trials, respectively, whereas those in Sample 2 received 30 and 10 reward for the 'rich' and 'lean' stimulus, respectively). Prior to analyses, we implemented a two-step procedure to identify outlier responses (Bogdan & Pizzagalli 2006; Pizzagalli *et al.* 2005). First, trials with reaction times (RT) less than 150 milliseconds or longer than 1500 milliseconds were excluded. Second, after removing outliers with step one, we naturally log transformed the remaining trials and calculated the RT mean

and standard deviation (SD) for each individual subject; trials that fell outside of the log-transformed mean ± 3 SD were excluded.

The main variable of interest was response bias, an empirically-based measure of reward learning, which measures the propensity to select a stimulus based on prior reinforcement history. Higher response bias values are reflective of a tendency to select the 'rich' stimulus as being displayed. Response bias was calculated according to the following formula:

$$\text{Response bias : } \log b = \frac{1}{2} \log \left(\frac{(\text{Rich}_{\text{correct}} + 0.5) * (\text{Lean}_{\text{incorrect}} + 0.5)}{(\text{Rich}_{\text{incorrect}} + 0.5) * (\text{Lean}_{\text{correct}} + 0.5)} \right)$$

This formula illustrates that increased response bias results from: (1) a high quantity of correct identifications of the 'rich' stimulus and misses for the 'lean' stimulus (i.e. incorrectly identifying the 'lean' stimulus as the rich stimulus) resulting in a large numerator, and (2) a low number of misses for the 'rich' stimulus and correct identifications of the 'lean' stimulus, resulting in a smaller denominator. The addition of 0.5 to each cell in this formula allows for the inclusion of data in which there were no incorrect responses.

To test the specificity of putative findings, control analyses were performed on discriminability, which provides a measure of the ability to discriminate between the two stimuli and is a measure of overall task performance or difficulty. Discriminability was calculated according to the following formula:

$$\text{Discriminability : } \log d = \frac{1}{2} \log \left(\frac{(\text{Rich}_{\text{correct}} + 0.5) * (\text{Lean}_{\text{correct}} + 0.5)}{(\text{Rich}_{\text{incorrect}} + 0.5) * (\text{Lean}_{\text{incorrect}} + 0.5)} \right)$$

Both measures, response bias and discriminability, were derived from the behavioral model of signal detection (Macmillan 2005).

Procedure

Sample 1

Participants completed two separate sessions. In the first session, the Structured Clinical Interview for the DSM-IV (SCID; First *et al.* 1997) was administered to ensure no past or current Axis I disorder was present (participants with past minor alcohol abuse, i.e. one symptom meeting threshold more than 2 years ago, were included, $n=2$). Eligible participants then completed a battery of questionnaires and provided a saliva sample for DNA analysis. During the second session participants performed the probabilistic reward task under a stress (threat-of-shock) and no-stress condition. In the stress condition, which was excluded from the present analyses, two electrodes were attached to the back of participants' right hand and participants were instructed that they would receive one to three electrical shocks during the stress condition and that the intensity of shocks would increase over time. For a complete description of this procedure please see Bogdan *et al.* (2011). The order of the stress and no stress condition was counterbalanced across participants. Only data from the no-stress condition was used for analyses.

Sample 2

Participants completed two separate sessions. In the first, the SCID (First *et al.* 1997) was administered to ensure that participants had no past or present Axis I disorders. Participants then completed several self-report measures assessing mood and stress, and provided a saliva sample for DNA analysis. In the second session, participants completed the probabilistic reward task. For a complete description of this procedure please see (Santesso *et al.* 2008).

Sample 3

Participants were recruited to complete the probabilistic reward learning task from a large ongoing study, the Duke Neurogenesis Study

(DNS), which assesses a wide range of behavioral, experiential and biological phenotypes among young-adult college students (Carey *et al.* 2015; Corral-Frías *et al.* 2015; Nikolova *et al.* 2014). Diagnosis of current DSM-IV Axis I and select Axis II disorders (Antisocial Personality Disorder and Borderline Personality Disorder) was assessed with the electronic Mini International Neuropsychiatric Interview (Sheehan *et al.* 1998) and SCID (First *et al.* 1997). These disorders were not exclusionary, as the DNS seeks to establish broad variability in multiple behavioral phenotypes related to psychopathology. After completing the initial portion of the study, some participants completed the probabilistic reward learning task on an additional day.

Genotyping

Samples 1 and 2

DNA obtained from saliva samples (OG-100; OG-25; Oragene; DNA Genotek Inc. 3000 – 500 Palladium Drive, Ottawa, Ontario, Canada K2V 1C2 DNA Genotek) was purified, extracted and hydrated; it was stored at -80°C when not in use. Primers were designed using Spectro DESIGNER v 2.0 software (Sequenom). Following a PCR, an iPLEX mass EXTEND reaction was performed. After baseline correction and peak identification, Sequenom SpectroTYPER v 2.0 software was used to analyze resulting spectra. Concordance for duplicate DNA in the current sample was 100%. *COMT* rs4680 did not deviate from Hardy-Weinberg equilibrium (HWE; all ethnicities: $\chi^2 = 0.372$, $P = 0.54$; European American sample only: $\chi^2 = 1.20$, $P = 0.27$; Sample 1: $\chi^2 = .055$, $P = 0.82$; Sample 2: $\chi^2 = 1.388$, $P = 0.24$).

Sample 3

DNA from participants within the DNS cohort was isolated from saliva derived from Oragene DNA self-collection kits (DNA Genotek) customized for 23andMe (www.23andme.com). DNA extraction and genotyping were performed by the National Genetics Institute (NGI), a CLIA-certified clinical laboratory and subsidiary of Laboratory Corporation of America. The Illumina HumanOmniExpress BeadChips and a custom array containing an additional ~300 000 SNPs were used to provide genome-wide data. *COMT* rs4680 did not deviate from HWE (all ethnicities: $\chi^2 = .118$, $P = 0.73$; European-American sample only: $\chi^2 = 2.44$, $P = 0.12$).

Data analysis

Because participants in each sample completed the same task with minor variations and response bias across these three samples did not differ, we combined samples. Kolmogorov-Smirnov test statistics indicated that the data did not significantly deviate from a normal distribution ($D = .04$; $P = 0.20$). As such, linear regressions (SPSS v.21) were used to test the association between *COMT* genotype and total response bias in the combined sample as well as in each sample individually. We used total response bias as our index of reward learning because this reflects the overall bias developed across the task. In addition, this metric has been previously associated with rs4680 genotype (Lancaster *et al.* 2012) and is robust (i.e. does not produce lower estimates) even in cases where individuals learn contingencies quickly. Given prior evidence that Met homozygotes have higher response bias relative to Val carriers, participants were separated into Val-allele carriers (Val/Val and Val/Met) and Met homozygotes (Met/Met) (Goetz *et al.* 2013; Lancaster *et al.* 2012, 2015). Additional results reporting an additive model are described in Appendix S1. Covariates included sex, study, and ethnicity (when applicable). Additionally, due to differences in age across samples and evidence that *COMT* enzyme activity differs according to age (Tunbridge *et al.* 2007), we also included age as a covariate. Finally, since some of the participants in our sample met criteria for one or more Axis I disorders (4.6%) according to a diagnostic interview (Tables S2 and S3), psychiatric diagnosis was also added as a covariate. Because the association between Val¹⁵⁸Met genotype and response bias has only been reported in European and European-American samples and due to population stratification concerns (Thomas & Witte 2002), primary analyses were conducted in European and European-American participants only ($n=208$). Supplemental analyses were conducted in the full population (see Appendix S1).

Meta-analysis

Literature search and analyses

We performed PubMed and Google scholar searches to identify *COMT* genotype and reward learning studies, using the probabilistic reward learning task of interest (Pizzagalli *et al.* 2005), published before December 2015. Search words included '*COMT* genotype', '*COMT* Val¹⁵⁸Met', 'rs4680', 'reward', 'reward learning', 'response bias' and 'probabilistic reward task'. This search yielded a total of three studies that were published between 2012 and 2015 (Goetz *et al.* 2013; Lancaster *et al.* 2012, 2015; Table 2). A weighted average for total response bias for Val carriers was calculated utilizing the means for Val/Val and Val/Met participants for a study that implemented an additive model (Goetz *et al.* 2013).

Analyses were performed using Revman 5.3 software (Cochrane IMS, Oxford, UK). The pooled effect was reported as a weighted mean difference (MD) with the corresponding 95% CI. Heterogeneity was assessed using I^2 and χ^2 tests, and a P value <0.10 was considered to be significant. Since heterogeneity was not present in this meta-analysis, the pooled effect size was calculated through a fixed-effects model. Forest plots were constructed with $P < 0.05$ considered to be significant.

Results

Response bias

COMT rs4680 (Val¹⁵⁸Met) genotype was significantly associated with total response bias ($\beta = 0.20$, $t = 2.75$, $P < 0.01$; $\Delta R^2 = 0.04$; Fig. 2) in the combined European-American samples ($n = 208$). Consistent with prior literature (Lancaster *et al.* 2012, 2015), Met allele homozygotes demonstrated relatively higher response bias ($M = 0.19$, $SD = 0.15$; $n = 55$) compared to Val allele carriers ($M = 0.12$, $SD = 0.16$; $n = 153$). The directionality of this relationship was also consistent when participants of all ethnicities were included in the analysis ($n = 303$); however, the effect of genotype was no longer significant ($\beta = 0.10$, $t = 1.60$, $P = 0.10$; Fig. S2). Although an additive genetic model showed consistent directional effects, these effects did not reach significance but showed a trending association (see Appendix S1 and Fig. S1).

The main effect of *COMT* rs4680 genotype on response bias was significant in the European-American population within the largest sample (Sample 2: $\beta = 0.19$, $t = 1.99$, $P = 0.04$; $n = 119$) but was not significant in Sample 3 ($\beta = 0.29$, $t = 1.50$; $P = 0.14$; $n = 33$) or 1 ($\beta = 0.15$, $t = 1.0$; $P = 0.32$; $n = 56$). Consistent with analyses combining data across samples, meta-analysis of all three independent samples from this study showed that Met-allele homozygotes had heightened response bias compared to Val carriers (MD: -0.07 ; 95% CI -0.12 to -0.02 ; $P < 0.01$; $n = 208$).

To ensure that our findings were specific to response bias and not due to differences in the ability to discriminate between the two different stimuli (i.e. discriminability), we conducted regression analyses using *COMT* rs4680 genotype as a predictor of discriminability. Highlighting the specificity of the response bias findings in this European American sample, *COMT* rs4680 genotype was not significantly associated with discriminability (Discriminability total: $\beta = -0.07$, $t = -1.60$, $P = 0.10$).

Meta-analysis

A pooled analysis of four studies (Goetz *et al.* 2013; Lancaster *et al.* 2012, 2015), present study, $n = 431$) of




Figure 2: *COMT* genotype associated with differences in total response bias in the probabilistic reward task. (a) In the European-American sample Met/Met participants ($n = 55$) demonstrated significantly greater total response bias than Val carriers ($n = 153$) ($\beta = .20$, $t = 2.75$, $P < 0.01$; $\Delta R^2 = .04$). Data points are jittered to allow for distribution visualization.


European/European-American participants revealed that response bias was significantly increased among Met-allele homozygotes ($n = 112$) compared to Val-allele carriers ($n = 319$; MD: -0.07 ; 95% CI -0.10 to -0.03 ; $P < 0.01$; Fig. 3). The test for heterogeneity was not significant ($I^2 = 48\%$; $P = 0.12$) confirming the appropriateness of a fixed effects model. Consistent with previous meta-analyses (Munafò *et al.* 2008), the effect size of the first published paper (Lancaster *et al.* 2012) was much larger than the effect size for the subsequent studies suggesting an overestimation of the effect in first published investigations (Ioannidis *et al.* 2001). Additionally, an analysis including participants from all ethnicities within the present sample ($n = 526$) revealed consistent results (MD: -0.04 ; 95% CI -0.08 to -0.003 ; $P < 0.05$; See Fig. S3).

Discussion

This study sought to replicate recently reported associations between *COMT* Val¹⁵⁸Met genotype and behavioral reward learning (Lancaster *et al.* 2012, 2015; Table 2). Consistent with these prior findings, our data suggest that individuals homozygous for the low enzymatic activity Met allele have relatively increased reward learning (as reflected by heightened response bias toward a stimulus more frequently associated with reward; Fig. 2). Further, a meta-analysis of four studies (Goetz *et al.* 2013; Lancaster *et al.* 2012, 2015 and the present study), using the same probabilistic reward learning task (Pizzagalli *et al.* 2005; $n = 431$), also produced a significant association between response bias and rs4680 genotype (Fig. 3). Consistent with our findings, recent complementary evidence suggests that *COMT* rs4680 genotype is

Table 2: Descriptive characteristics of studies included in meta-analysis

Study	Year	Age	Sex (% female)	Ancestry	Country of origin	Allele frequency			Total N
						Met/Met	Val/Met	Val/Val	
Lancaster <i>et al.</i>	2012	22.7 ± 4.2	61.42%	European	England	19	25	26	70
Goetz <i>et al.</i>	2013	21.3 ± 2.7	59.32%	European-American	United States	14	28	17	59
Lancaster <i>et al.</i>	2015	22.2 ± 4.6	43.56%	European	England	24	54	20	98
Current manuscript		21.8 ± 3.9	64.90%	European-American	United States	55	97	56	208

**Figure 3: Forest plot of the pooled effect of COMT genotype on total response bias.** Size of square is proportional to sample size. CI: confidence interval; df: degrees of freedom; IV: Inverse Variance (statistical method). Lancaster *et al.* (2015) report a significant *COMT* rs4680 genotype x block interaction; here we depict the effect for the main effect of *COMT* rs4680 genotype on response bias.

also associated with other aspects of reward function, including positive affect in response to rewarding experiences (Wichers *et al.* 2007) and reward seeking behavior (Lancaster *et al.* 2012). Collectively, these findings across studies provide evidence for an association between *COMT* rs4680 genotype and individual differences in reward processing, which may in turn, confer variability in vulnerability to a host of psychopathologies.

Putative neural mechanisms

While this study did not examine putative neural mechanisms through which *COMT* rs4680 genotype may be associated with individual differences in reward learning, emerging literature probing associations between *COMT* rs4680 genotype and neural phenotypes allows for informed speculation. This research suggests differential DA function associated with *COMT* rs4680 genotype wherein Met-allele homozygosity results in higher PFC DA levels relative to Val-allele homozygosity due to 40% fold reduction in *COMT* activity (Chen *et al.* 2004). This Met-allele driven variability in PFC DA may influence phasic reward prediction signals, stimulus signal-to-noise ratios as well as working memory to produce differences in reward learning. However, as noted below, each of these interpretations is also challenged by conflicting evidence.

Phasic changes in subcortical DA neuron firing are thought to encode differences in reward prediction errors (i.e. difference between expected and observed value), which are crucial signals for reward learning (Bayer & Glimcher 2005; Garris *et al.* 1999; Schultz 2002, 2007). The expression of *COMT* in subcortical regions is minimal and the direct effect on subcortical DA cell activity is unknown. However,

midbrain and striatal dopaminergic neurons are regulated by the PFC (Seamans & Yang 2004), and the *COMT* Val allele is associated with increased tyrosine hydroxylase expression within the midbrain and, hence, presumably increased DA synthesis (Akil *et al.* 2003; Meyer-Lindenberg *et al.* 2005). This putative increase in DA synthesis may decrease the ability to detect phasic activity necessary for reward prediction, leading to decreased reward learning in Val-allele carriers (Pizzagalli *et al.* 2008; Santesso *et al.* 2009). Moreover, rs4680 genotype has been associated with individual differences in the functional interactions of subcortical and prefrontal regions during a working memory task (Meyer-Lindenberg *et al.* 2005) suggesting that *COMT* rs4680 genotype-related differences in PFC-subcortical interactions may contribute to reward learning.

Notably however, while this interpretation is consistent with our understanding of the role of subcortical DA and reward learning, direct neuroimaging studies of *COMT* rs4680 genotype associations with reward-related brain activation, which is believed to be tied to DA signaling (Knutson & Gibbs 2007), have yielded conflicting evidence (Antypa *et al.* 2013; Camara *et al.* 2009; Dreher *et al.* 2009; Forbes *et al.* 2009; Schmack *et al.* 2008; Yacubian *et al.* 2007). For instance, some studies have shown increased reward-related ventral striatum reactivity in Met homozygotes (Dreher *et al.* 2009; Schmack *et al.* 2008; Yacubian *et al.* 2007) while others have shown increased activation in Val homozygotes (Camara *et al.* 2009). While these contradictory findings call into question the potential impact of *COMT* genotype on reward function through its effect on subcortical DA, it is important to note that none of these tasks were designed to evaluate reward learning specifically (as opposed to other forms of reward processing such as anticipating or receiving money).

It is also possible that false positive associations (Farrell *et al.* 2015; Lee & Song 2015; Munafò *et al.* 2005; Nickl-Jockschat *et al.* 2015) may contribute to these equivocal results.

Alternatively, though not mutually exclusive, the effects of rs4680 genotype on reward learning may arise from its effects on prefrontal DA function and related behaviors. In addition to striatal reward prediction errors, a wide variety of higher order brain function, including executive control and working memory, likely contributes to reward learning (Collins & Frank 2012). Extensive working memory research suggests COMT-related effects on prefrontal DA may modulate signal-to-noise ratio allowing task-related information to be prioritized, potentially facilitating learning of novel stimulus-reward pairings (Akil *et al.* 2003; Meyer-Lindenberg & Weinberger 2006; Seamans & Yang 2004 but see also Nickl-Jockschat *et al.* 2015). Further, this literature suggests that in the context of working memory, an optimum level of DA stimulation is necessary to reach the highest signal-to-noise ratio, placing Met homozygotes at the height of this inverted u-curve (Meyer-Lindenberg *et al.* 2005). Moreover, behavioral and *in silico* experiments suggest that prefrontal function may contribute to reward learning by influencing initial learning acquisition rate (Collins & Frank 2012). Supporting this interpretation, genetic association studies have linked polymorphisms associated with variability in subcortical DA signaling (e.g. *DARPP-32* and *DRD2*) to individual differences in learning rates after initial learning has occurred, and polymorphisms associated with cortical DA function (e.g. *COMT*) with learning rates during initial acquisition (Frank *et al.* 2007). Thus, although the effects of *COMT* rs4680 genotype on subcortical function have been hypothesized (Bildner *et al.* 2004), evidence (Huotari *et al.* 2002) suggests that an explanation based on prefrontal regulation of striatal DA metabolism via top-down projections may also be important (Matsumoto *et al.* 2003).

While this interpretation could potentially account for the behavioral effects observed here, it is challenged by recent meta-analyses suggesting that *COMT* rs4680 genotype may have no main effect on higher order executive function such as working memory (Nickl-Jockschat *et al.* 2015). Notably, it is possible that Val allele-specific patterns of methylation may contribute to this contradictory literature, as the Val-allele homozygotes have a CpG methylation site that Met-allele carriers do not. Moreover, methylation at this site is related to stress exposure and variability in behavioral and neural working memory phenotypes (Ursini *et al.* 2011). Specifically, Val-allele homozygotes with low stress levels and heightened methylation in this region have working memory-related neural function and behavior comparable to Met-allele carriers highlighting the importance of considering methylation and stress in future *COMT* rs4680 genotype research (Ursini *et al.* 2011).

Vulnerability to psychopathology

Recent theoretical and empirical evidence suggests that reward processing deficits within psychiatric disorders may be closely linked to motivation, reward learning and reward decision making, rather than hedonic response (Barch *et al.* 2015; Pizzagalli 2014). Since positive reinforcement increases

the likelihood of behaviors linked to them, reward learning dysfunction may reduce motivation to pursue rewards, thus increasing the probability of symptom persistence or even exacerbation of psychopathology (Pizzagalli 2014). Consistent with this hypothesis, behavioral reward learning, as measured by the task used in this study, has been associated with anhedonic symptoms and depression (Luking *et al.* 2015b,2015a; Pizzagalli *et al.* 2005) as well as chronicity of symptoms after antidepressant treatment (Vrieze *et al.* 2013). Accordingly, reward learning deficits observed in *COMT* rs4680 Val-allele carriers may place them at greater risk for psychopathology characterized by deficient reward processing as has been reported in some (Baune *et al.* 2008; Benedetti *et al.* 2009, 2010; Spronk *et al.* 2011; Yoshida *et al.* 2008), but not all (Szegedi *et al.* 2005), studies.

Limitations and conclusions

Interpretation of the current results should be considered in the context of study limitations. First, our sample (even the pooled meta-analytic data) is small for a genetic association study making our estimated effect imprecise. Second, our results were only significant when all subsamples were combined (or meta-analyzed), and in our largest dataset (Sample 2). The relationship was not significant in our other samples, though it showed a similar directional effect across all. The non-significant results from Sample 1 may be partially attributable to the original design of the study. Here participants performed the probabilistic reward learning task under stress and no-stress conditions, where the order was counterbalanced across subjects (Bogdan *et al.* 2010, 2011). While this study did not use data from the stress condition, given preliminary evidence of Gene x Environment interactions at this locus (Craddock *et al.* 2006), it is possible that this study design and the presence of the stress condition weakened the link between *COMT* genotype and reward learning. Moreover, our three samples differed in sex, ethnicity and age distribution as well as the version of the probabilistic reward task used. These study-related differences may have added variability to our reported effects. In an attempt to account for this possibility, study differences including, the study of origin, sex and age, were included as covariates in our analysis. Further, we analyzed each sample independently and conducted a meta-analysis, which resulted in the same conclusion.

Third, results were only significant when analyses were constrained to European-American individuals. However, the directionality of the effect was consistent when all ethnicities were included in analyses (Fig. S2) and a meta-analysis across published studies (including the present data) also yielded evidence of significant association when including individuals of all ancestral origins (Fig. S3). It is important to note that this is the only study to date to contain a sample of mixed ethnicities so the role of ancestral origin in *COMT* Val¹⁵⁸Met genotype-response bias phenotype associations is unclear. Notably, among other phenotypes, there is evidence of differential association according to ancestry (e.g. Domschke *et al.* 2007; Hosak 2007).

Fourth, our meta-analysis only included data from previously published research. In light of publication bias for positive as opposed to null findings (Hirschhorn *et al.* 2002; Munafo *et al.* 2004), it is possible that additional data are available which do not report the associations described herein and may have led to a biased meta-analysis. Along with these previous reports, our study thus highlights the importance of not only replicating genetic association studies but also performing meta-analyses in an attempt to more accurately measure effect sizes (Munafo *et al.* 2008). Lastly, meta-analyses were conducted using a fixed effects model. While the heterogeneity observed in the data support such a model, the larger studies included within the meta-analysis by definition contributes more to the weighted average. Notably, a random effects model showed trending effects ($P = .08$) in the same direction as the fixed effects model.

These limitations notwithstanding, the present study suggests that a common genetic variant within the *COMT* gene (rs4680) is associated with individual differences in reward learning. Our study further highlights the importance of replication and meta-analyses in genetic association studies. While these findings shed light on how this functional genetic polymorphism is important in the appearance of individual differences in reward learning, further research is needed to elucidate the potential neural mechanisms underlying these behavioral associations and to trace such associations to the development of psychopathology.

References

- Akil, M., Kolachana, B.S., Rothmond, D.A., Hyde, T.M., Weinberger, D.R. & Kleinman, J.E. (2003) Catechol-O-methyltransferase genotype and dopamine regulation in the human brain. *J Neurosci* **23**, 2008–2013.
- Antypa, N., Drago, A. & Serretti, A. (2013) The role of COMT gene variants in depression: bridging neuropsychological, behavioral and clinical phenotypes. *Neurosci Biobehav Rev* **37**, 1597–1610.
- Barch, D.M., Pagliaccio, D. & Luking, K. (2015) Mechanisms underlying motivational deficits in psychopathology: similarities and differences in depression and schizophrenia. *Curr Top Behav Neurosci*. DOI: 10.1007/7854_2015_376.
- Baune, B.T., Hohoff, C., Berger, K., Neumann, A., Mortensen, S., Roehrs, T., Deckert, J., Arolt, V. & Domschke, K. (2008) Association of the COMT val¹⁵⁸met variant with antidepressant treatment response in major depression. *Neuropsychopharmacology* **33**, 924–932.
- Bayer, H.M. & Glimcher, P.W. (2005) Midbrain dopamine neurons encode a quantitative reward prediction error signal. *Neuron* **47**, 129–141.
- Benedetti, F., Colombo, C., Pirovano, A., Marino, E. & Smeraldi, E. (2009) The catechol-O-methyltransferase Val^(108/158)Met polymorphism affects antidepressant response to paroxetine in a naturalistic setting. *Psychopharmacology (Berl)* **203**, 155–160.
- Benedetti, F., Dallaspesza, S., Colombo, C., Lorenzi, C., Pirovano, A. & Smeraldi, E. (2010) Effect of catechol-O-methyltransferase Val^(108/158)Met polymorphism on antidepressant efficacy of fluvoxamine. *Eur Psychiatry* **25**, 476–478.
- Bilder, R.M., Volavka, J., Lachman, H.M. & Grace, A.A. (2004) The catechol-O-methyltransferase polymorphism: relations to the tonic-phasic dopamine hypothesis and neuropsychiatric phenotypes. *Neuropsychopharmacology* **29**, 1943–1961.
- Bogdan, R. & Pizzagalli, D.A. (2006) Acute stress reduces reward responsiveness: implications for depression. *Biol Psychiatry* **60**, 1147–1154.
- Bogdan, R. & Pizzagalli, D.A. (2009) The heritability of hedonic capacity and perceived stress: a twin study evaluation of candidate depressive phenotypes. *Psychol Med* **39**, 211–218.
- Bogdan, R., Perlis, R.H., Fagerness, J. & Pizzagalli, D.A. (2010) The impact of mineralocorticoid receptor ISO/VAL genotype (rs5522) and stress on reward learning. *Genes Brain Behav* **9**, 658–667.
- Bogdan, R., Santesso, D.L., Fagerness, J., Perlis, R.H. & Pizzagalli, D.A. (2011) Corticotropin-releasing hormone receptor type 1 (CRHR1) genetic variation and stress interact to influence reward learning. *J Neurosci* **31**, 13246–13254.
- Bogdan, R., Nikolova, Y.S. & Pizzagalli, D.A. (2013) Neurogenetics of depression: a focus on reward processing and stress sensitivity. *Neurobiol Dis* **52**, 12–23.
- Buckholz, J.W. & Meyer-Lindenberg, A. (2012) Psychopathology and the human connectome: toward a transdiagnostic model of risk for mental illness. *Neuron* **74**, 990–1004.
- Camara, E., Kramer, U.M., Cunillera, T., Marco-Pallares, J., Cucurell, D., Nager, W., Mestres-Misse, A., Bauer, P., Schule, R., Schols, L., Tempelmann, C., Rodriguez-Fornells, A. & Munte, T.F. (2009) The effects of COMT (Val^{108/158}Met) and DRD4 (SNP-521) dopamine genotypes on brain activations related to valence and magnitude of rewards. *Cereb Cortex* **20**, 1985–1996.
- Carey, C.E., Agrawal, A., Zhang, B., Conley, E.D., Degenhardt, L., Heath, A.C., Li, D., Lynskey, M.T., Martin, N.G., Montgomery, G.W., Wang, T., Bierut, L.J., Hariri, A.R., Nelson, E.C. & Bogdan, R. (2015) Monoacylglycerol lipase (MGLL) polymorphism rs604300 interacts with childhood adversity to predict cannabis dependence symptoms and amygdala habituation: evidence from an endocannabinoid system-level analysis. *J Abnorm Psychol* **124**, 860–877.
- Chen, J., Lipska, B.K., Halim, N., Ma, Q.D., Matsumoto, M., Melhem, S., Kolachana, B.S., Hyde, T.M., Herman, M.M., Apud, J., Egan, M.F., Kleinman, J.E. & Weinberger, D.R. (2004) Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): effects on mRNA, protein, and enzyme activity in postmortem human brain. *Am J Hum Genet* **75**, 807–821.
- Collins, A.G. & Frank, M.J. (2012) How much of reinforcement learning is working memory, not reinforcement learning? A behavioral, computational, and neurogenetic analysis. *Eur J Neurosci* **35**, 1024–1035.
- Corral-Frías, N.S., Nikolova, Y.S., Michalski, L.J., Baranger, D.A., Hariri, A.R. & Bogdan, R. (2015) Stress-related anhedonia is associated with ventral striatum reactivity to reward and transdiagnostic psychiatric symptomatology. *Psychol Med* **45**, 2605–2617.
- Craddock, N., Owen, M.J. & O'Donovan, M.C. (2006) The catechol-O-methyl transferase (COMT) gene as a candidate for psychiatric phenotypes: evidence and lessons. *Molecular Psychiatry* **11**, 446–458.
- Der-Avakian, A., D'Souza, M.S., Pizzagalli, D.A. & Markou, A. (2013) Assessment of reward responsiveness in the response bias probabilistic reward task in rats: implications for cross-species translational research. *Transl Psychiatry* **3**, e297.
- Docherty, A.R. & Sponheim, S.R. (2008) Anhedonia as a phenotype for the Val¹⁵⁸Met COMT polymorphism in relatives of patients with schizophrenia. *J Abnorm Psychol* **117**, 788–798.
- Domschke, K., Deckert, J., O'Donovan, M.C. & Glatt, S.J. (2007) Meta-analysis of COMT val¹⁵⁸met in panic disorder: ethnic heterogeneity and gender specificity. *Am J Med Genet B Neuropsychiatr Genet* **144B**, 667–673.
- Dreher, J.C., Kohn, P., Kolachana, B., Weinberger, D.R. & Berman, K.F. (2009) Variation in dopamine genes influences responsivity of the human reward system. *Proc Natl Acad Sci U S A* **106**, 617–622.
- Duncan, L.E. & Keller, M.C. (2011) A critical review of the first 10 years of candidate gene-by-environment interaction research in psychiatry. *Am J Psychiatry* **168**, 1041–1049.
- Farrell, M.S., Werge, T., Sklar, P., Owen, M.J., Ophoff, R.A., O'Donovan, M.C., Corvin, A., Cichon, S. & Sullivan, P.F. (2015) Evaluating historical candidate genes for schizophrenia. *Mol Psychiatry* **20**, 555–562.

- First, M.B., Gibbon, M., Spitzer, R.L., Williams, J.B.W. & Benjamin, L.S. (1997) *Structured Clinical Interview for DSM-IV Axis II Personality Disorders, (SCID-II)*. American Psychiatric Press, Inc., Washington, DC.
- Forbes, E.E., Brown, S.M., Kimak, M., Ferrell, R.E., Manuck, S.B. & Hariri, A.R. (2009) Genetic variation in components of dopamine neurotransmission impacts ventral striatal reactivity associated with impulsivity. *Mol Psychiatry* **14**, 60–70.
- Frank, M.J., Moustafa, A.A., Haughey, H.M., Curran, T. & Hutchison, K.E. (2007) Genetic triple dissociation reveals multiple roles for dopamine in reinforcement learning. *Proc Natl Acad Sci U S A* **104**, 16311–16316.
- Garfield, J.B.B., Lubman, D.I. & Yucel, M. (2014) Anhedonia in substance use disorders: a systematic review of its nature, course and clinical correlates. *Aust N Z J Psychiatry* **48**, 36–51.
- Garris, P.A., Kilpatrick, M., Bunin, M.A., Michael, D., Walker, Q.D. & Wightman, R.M. (1999) Dissociation of dopamine release in the nucleus accumbens from intracranial self-stimulation. *Nature* **398**, 67–69.
- Gatt, J.M., Burton, K.L., Williams, L.M. & Schofield, P.R. (2015) Specific and common genes implicated across major mental disorders: a review of meta-analysis studies. *J Psychiatr Res* **60**, 1–13.
- Goetz, E.L., Hariri, A.R., Pizzagalli, D.A. & Strauman, T.J. (2013) Genetic moderation of the association between regulatory focus and reward responsiveness: a proof-of-concept study. *Biol Mood Anxiety Disord* **3**, 3.
- Gorwood, P. (2008) Neurobiological mechanisms of anhedonia. *Dialogues Clin Neurosci* **10**, 291–299.
- Hatzigiakoumis, D.S., Martinotti, G., Giannantonio, M.D. & Janiri, L. (2011) Anhedonia and substance dependence: clinical correlates and treatment options. *Front Psychiatry* **2**, 10.
- Herrnstein, R.J. (1961) Relative and absolute strength of response as a function of frequency of reinforcement. *J Exp Anal Behav* **4**, 267–272.
- Hirschhorn, J.N., Lohmueller, K., Byrne, E. & Hirschhorn, K. (2002) A comprehensive review of genetic association studies. *Genet Med* **4**, 45–61.
- Hosak, L. (2007) Role of the COMT gene Val¹⁵⁸Met polymorphism in mental disorders: a review. *Eur Psychiatry* **22**, 276–281.
- Huotari, M., Santha, M., Lucas, L.R., Karayiorgou, M., Gogos, J.A. & Mannisto, P.T. (2002) Effect of dopamine uptake inhibition on brain catecholamine levels and locomotion in catechol-O-methyltransferase-disrupted mice. *J Pharmacol Exp Ther* **303**, 1309–1316.
- Huys, Q.J., Pizzagalli, D.A., Bogdan, R. & Dayan, P. (2013) Mapping anhedonia onto reinforcement learning: a behavioural meta-analysis. *Biol Mood Anxiety Disord* **3**, 12.
- Ioannidis, J.P., Ntzani, E.E., Trikalinos, T.A. & Contopoulos-Ioannidis, D.G. (2001) Replication validity of genetic association studies. *Nat Genet* **29**, 306–309.
- Knutson, B. & Gibbs, S.E. (2007) Linking nucleus accumbens dopamine and blood oxygenation. *Psychopharmacology (Berl)* **191**, 813–822.
- Lancaster, T.M., Linden, D.E. & Heerey, E.A. (2012) COMT val¹⁵⁸met predicts reward responsiveness in humans. *Genes Brain Behav* **11**, 986–992.
- Lancaster, T.M., Heerey, E.A., Mantripragada, K. & Linden, D.E. (2015) Replication study implicates COMT val¹⁵⁸met polymorphism as a modulator of probabilistic reward learning. *Genes Brain Behav* **14**, 486–492.
- Lauwereyns, J., Watanabe, K., Coe, B. & Hikosaka, O. (2002) A neural correlate of response bias in monkey caudate nucleus. *Nature* **418**, 413–417.
- Lee, L.O. & Prescott, C.A. (2014) Association of the catechol-O-methyltransferase val¹⁵⁸met polymorphism and anxiety-related traits: a meta-analysis. *Psychiatr Genet* **24**, 52–69.
- Lee, Y.H. & Song, G.G. (2015) BDNF 196G/A and COMT Val¹⁵⁸Met polymorphisms and susceptibility to ADHD: a meta-analysis. *J Atten Disord*. DOI: 10.1177/1087054715570389.
- Liverant, G.I., Sloan, D.M., Pizzagalli, D.A., Harte, C.B., Kamholz, B.W., Rosebrock, L.E., Cohen, A.L., Fava, M. & Kaplan, G.B. (2014) Associations among smoking, anhedonia, and reward learning in depression. *Behav Ther* **45**, 651–663.
- Loas, G. (1996) Vulnerability to depression: a model centered on anhedonia. *J Affect Disord* **41**, 39–53.
- Luking, K.R., Neiman, J.S., Luby, J.L. & Barch, D.M. (2015a) Reduced hedonic capacity/approach motivation relates to blunted responsiveness to gain and loss feedback in children. *J Clin Child Adolesc Psychol*, 1–13. DOI:10.1080/15374416.2015.1012721.
- Luking, K.R., Pagliaccio, D., Luby, J.L. & Barch, D.M. (2015b) Child gain approach and loss avoidance behavior: relationships with depression risk, negative mood, and anhedonia. *J Am Acad Child Adolesc Psychiatry* **54**, 643–651.
- Macmillan, N.A., (2005) *Detection theory: a user's guide*, 2nd ed. Lawrence Erlbaum Associates, Mahwah, N.J.
- Mannisto, P.T. & Kaakkola, S. (1999) Catechol-O-methyltransferase (COMT): biochemistry, molecular biology, pharmacology, and clinical efficacy of the new selective COMT inhibitors. *Pharmacol Rev* **51**, 593–628.
- Matsumoto, M., Weickert, C.S., Akil, M., Lipska, B.K., Hyde, T.M., Herman, M.M., Kleinman, J.E. & Weinberger, D.R. (2003) Catechol O-methyltransferase mRNA expression in human and rat brain: evidence for a role in cortical neuronal function. *Neuroscience* **116**, 127–137.
- Meehl, P.E. (1975) Hedonic capacity: some conjectures. *Bull Menninger Clin* **39**, 295–307.
- Meyer-Lindenberg, A. & Weinberger, D.R. (2006) Intermediate phenotypes and genetic mechanisms of psychiatric disorders. *Nat Rev Neurosci* **7**, 818–827.
- Meyer-Lindenberg, A., Kohn, P.D., Kolachana, B., Kippenhan, S., McInerney-Leo, A., Nussbaum, R., Weinberger, D.R. & Berman, K.F. (2005) Midbrain dopamine and prefrontal function in humans: interaction and modulation by COMT genotype. *Nat Neurosci* **8**, 594–596.
- Morris, B.H. & Rottenberg, J. (2015) Heightened reward learning under stress in generalized anxiety disorder: a predictor of depression resistance? *J Abnorm Psychol* **124**, 115–127.
- Munafò, M.R., Clark, T.G. & Flint, J. (2004) Assessing publication bias in genetic association studies: evidence from a recent meta-analysis. *Psychiatry Res* **129**, 39–44.
- Munafò, M.R., Bowes, L., Clark, T.G. & Flint, J. (2005) Lack of association of the COMT (Val¹⁵⁸/108Met) gene and schizophrenia: a meta-analysis of case-control studies. *Mol Psychiatry* **10**, 765–770.
- Munafò, M.R., Brown, S.M. & Hariri, A.R. (2008) Serotonin transporter (5-HTTLPR) genotype and amygdala activation: a meta-analysis. *Biol Psychiatry* **63**, 852–857.
- Nickl-Jockschat, T., Janouschek, H., Eickhoff, S.B. & Eickhoff, C.R. (2015) Lack of meta-analytic evidence for an impact of COMT Val¹⁵⁸Met genotype on brain activation during working memory tasks. *Biol Psychiatry* **78**, e43–46.
- Nikolova, Y.S., Ferrell, R.E., Manuck, S.B. & Hariri, A.R. (2011) Multilocus genetic profile for dopamine signaling predicts ventral striatum reactivity. *Neuropsychopharmacology* **36**, 1940–1947.
- Nikolova, Y.S., Bogdan, R., Brigidi, B.D. & Hariri, A.R. (2012) Ventral striatum reactivity to reward and recent life stress interact to predict positive affect. *Biol Psychiatry* **72**, 157–163.
- Nikolova, Y.S., Koenen, K.C., Galea, S., Wang, C.M., Seney, M.L., Sibille, E., Williamson, D.E. & Hariri, A.R. (2014) Beyond genotype: serotonin transporter epigenetic modification predicts human brain function. *Nat Neurosci* **17**, 1153–1155.
- Pergadia, M.L., Der-Avakian, A., D'Souza, M.S., Madden, P.A., Heath, A.C., Shiffman, S., Markou, A. & Pizzagalli, D.A. (2014) Association between nicotine withdrawal and reward responsiveness in humans and rats. *JAMA Psychiatry* **71**, 1238–1245.
- Pizzagalli, D.A. (2014) Depression, stress, and anhedonia: toward a synthesis and integrated model. *Ann Rev Clin Psychol* **10**, 393–423.

- Pizzagalli, D.A., Jahn, A.L. & O'Shea, J.P. (2005) Toward an objective characterization of an anhedonic phenotype: a signal-detection approach. *Biol Psychiatry* **57**, 319–327.
- Pizzagalli, D.A., Evins, A.E., Schetter, E.C., Frank, M.J., Pajtas, P.E., Santesso, D.L. & Culhane, M. (2008) Single dose of a dopamine agonist impairs reinforcement learning in humans: behavioral evidence from a laboratory-based measure of reward responsiveness. *Psychopharmacology (Berl)* **196**, 221–232.
- Plomin, R., DeFries, J.C., Knopik, V.S. & Neiderhiser, J.M. (2016) Top 10 replicated findings from behavioral genetics. *Perspect Psychol Sci* **11**, 3–23.
- Santesso, D.L., Steele, K.T., Bogdan, R., Holmes, A.J., Deveney, C.M., Meites, T.M. & Pizzagalli, D.A. (2008) Enhanced negative feedback responses in remitted depression. *Neuroreport* **19**, 1045–1048.
- Santesso, D.L., Evins, A.E., Frank, M.J., Schetter, E.C., Bogdan, R. & Pizzagalli, D.A. (2009) Single dose of a dopamine agonist impairs reinforcement learning in humans: evidence from event-related potentials and computational modeling of striatal-cortical function. *Hum Brain Mapp* **30**, 1963–1976.
- Schmack, K., Schlagenhaut, F., Sterzer, P., Wrase, J., Beck, A., Demberler, T., Kalus, P., Puls, I., Sander, T., Heinz, A. & Gallinat, J. (2008) Catechol-O-methyltransferase val¹⁵⁸met genotype influences neural processing of reward anticipation. *Neuroimage* **42**, 1631–1638.
- Schultz, W. (2002) Getting formal with dopamine and reward. *Neuron* **36**, 241–263.
- Schultz, W. (2007) Behavioral dopamine signals. *Trends Neurosci* **30**, 203–210.
- Schultz, W. (2015) Neuronal reward and decision signals: from theories to data. *Physiol Rev* **95**, 853–951.
- Scornaiencki, R., Cantrup, R., Rushlow, W.J. & Rajakumar, N. (2009) Prefrontal cortical D1 dopamine receptors modulate subcortical D2 dopamine receptor-mediated stress responsiveness. *Int J Neuropsychopharmacol* **12**, 1195–1208.
- Seamans, J.K. & Yang, C.R. (2004) The principal features and mechanisms of dopamine modulation in the prefrontal cortex. *Prog Neurobiol* **74**, 1–58.
- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R. & Dunbar, G.C. (1998) The mini-international neuropsychiatric interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* **59** (Suppl 20), 22–33quiz 34–57.
- Slifstein, M., Kolachana, B., Simpson, E.H., Tabares, P., Cheng, B., Duvall, M., Frankle, W.G., Weinberger, D.R., Laruelle, M. & Abi-Dargham, A. (2008) COMT genotype predicts cortical-limbic D1 receptor availability measured with [¹¹C]NNC112 and PET. *Mol Psychiatry* **13**, 821–827.
- Spronk, D., Arns, M., Barnett, K.J., Cooper, N.J. & Gordon, E. (2011) An investigation of EEG, genetic and cognitive markers of treatment response to antidepressant medication in patients with major depressive disorder: a pilot study. *J Affect Disord* **128**, 41–48.
- Szegedi, A., Rujescu, D., Tadic, A., Muller, M.J., Kohnen, R., Stassen, H.H. & Dahmen, N. (2005) The catechol-O-methyltransferase Val^{108/158}Met polymorphism affects short-term treatment response to mirtazapine, but not to paroxetine in major depression. *Pharmacogenomics J* **5**, 49–53.
- Thomas, D.C. & Witte, J.S. (2002) Point: population stratification: a problem for case-control studies of candidate-gene associations? *Cancer Epidemiol Biomarkers Prev* **11**, 505–512.
- Tripp, G. & Alsop, B. (1999) Sensitivity to reward frequency in boys with attention deficit hyperactivity disorder. *J Clin Child Psychol* **28**, 366–375.
- Tunbridge, E.M., Bannerman, D.M., Sharp, T. & Harrison, P.J. (2004) Catechol-o-methyltransferase inhibition improves set-shifting performance and elevates stimulated dopamine release in the rat prefrontal cortex. *J Neurosci* **24**, 5331–5335.
- Tunbridge, E.M., Weickert, C.S., Kleinman, J.E., Herman, M.M., Chen, J., Kolachana, B.S., Harrison, P.J. & Weinberger, D.R. (2007) Catechol-o-methyltransferase enzyme activity and protein expression in human prefrontal cortex across the postnatal lifespan. *Cereb Cortex* **17**, 1206–1212.
- Ursini, G., Bollati, V., Fazio, L., Porcelli, A., Iacovelli, L., Catalani, A., Sinibaldi, L., Gelao, B., Romano, R., Rampino, A., Taurisano, P., Mancini, M., Di Giorgio, A., Popolizio, T., Baccarelli, A., De Blasi, A., Blasi, G. & Bertolino, A. (2011) Stress-related methylation of the catechol-O-methyltransferase Val¹⁵⁸ allele predicts human prefrontal cognition and activity. *J Neurosci* **31**, 6692–6698.
- Vrieze, E., Pizzagalli, D.A., Demyttenaere, K., Hompes, T., Siennaert, P., de Boer, P., Schmidt, M. & Claes, S. (2013) Reduced reward learning predicts outcome in major depressive disorder. *Biol Psychiatry* **73**, 639–645.
- Wichers, M., Aguilera, M., Kenis, G., Krabbendam, L., Myin-Germeys, I., Jacobs, N., Peeters, F., Derom, C., Vlietinck, R., Mengelers, R., Delespaul, P. & van Os, J. (2007) The catechol-O-methyl transferase val¹⁵⁸met polymorphism and experience of reward in the flow of daily life. *Neuropsychopharmacology* **33**, 3030–3036.
- Yacubian, J., Sommer, T., Schroeder, K., Glascher, J., Kalisch, R., Leuenberger, B., Braus, D.F. & Buchel, C. (2007) Gene-gene interaction associated with neural reward sensitivity. *Proc Natl Acad Sci U S A* **104**, 8125–8130.
- Yoshida, K., Higuchi, H., Takahashi, H., Kamata, M., Sato, K., Inoue, K., Suzuki, T., Itoh, K. & Ozaki, N. (2008) Influence of the tyrosine hydroxylase val⁸¹met polymorphism and catechol-O-methyltransferase val¹⁵⁸met polymorphism on the antidepressant effect of milnacipran. *Hum Psychopharmacol* **23**, 121–128.

Acknowledgments

We are thankful to the Duke Neurogenetics Study (DNS) and Center for Depression, Anxiety and Stress Research (CDASR) participants and staff. Studies conducted at Harvard University were supported by NIMH Grant R01-MH068376 (awarded to D.A.P.), and a Sackler Scholar in Psychobiology (awarded to R.B.). D.A.P. was also partially supported by NIMH grant MH101521. The Duke Neurogenetics Study is supported by Duke University and the National Institutes of Health (NIDA R01-DA033369). A.R.H. receives additional support from the National Institutes of Health (NIDA R01-DA031579). N.C.S. was supported by NIMH (T32-MH014677). R.B. was supported by the Klingenstein Third Generation Foundation and receives additional support from the National Institutes of Health (NIA R01-AG045231). L.J.M. was supported by the National Institutes of Health (T32-GM081739). In the past three years, Dr. Pizzagalli has received honoraria/consulting fees from Otsuka America Pharmaceutical and Pfizer for activities unrelated to this project. All other authors report no conflict of interest.

Supporting Information

Additional supporting information may be found in the online version of this article at the publisher's web-site:

Figure S1: Additive model of *COMT* genotype and response bias. Results demonstrated a non-significant but trending association between *COMT* genotype and response bias in the additive model ($\beta = .12$, $t = 1.76$, $P = 0.07$; $n = 208$). Data points are jittered to allow for distribution visualization.

Figure S2: *COMT* genotype and response bias in whole sample (all ethnicities). Results demonstrated a non-significant association between *COMT* genotype and response bias when all ethnicities were included ($\beta = .10$,

$t = 1.60$, $P = 0.10$, $n = 303$). Data points are jittered to allow for distribution visualization.

Figure S3: Forest plot of the pooled effect of *COMT* genotype on response bias (all ethnicities included). Size of square is proportional to sample size. CI: confidence interval; df: degrees of freedom; IV: Inverse Variance (statistical method) RB: response bias. Lancaster *et al.* (2015) report a significant *COMT* rs4680 genotype \times block interaction; here we depict the effect for the main effect of *COMT* rs4680 genotype on response bias.

Figure S4: Forest plot of the pooled effect of *COMT* genotype on response bias of each sample in the study. Size of square is proportional to sample size. CI: confidence interval; df: degrees of freedom; IV: Inverse Variance (statistical method) RB: response bias.

Table S1: Sample demographics across all ethnicities.

Table S2: DSM-IV Axis I Disorder Diagnosis in all ethnicities.

Table S3: DSM-IV Axis I Disorder Diagnosis in European American participants.